

FEDERAL PUBLIC SERVICE COMMISSION



COMPETITIVE EXAMINATION FOR RECRUITMENT TO POSTS IN BS-17 UNDER THE FEDERAL GOVERNMENT, 2012

Roll Number

APPLIED MATHS, PAPER-II

TIME ALLOWED: THREE HOURS

MAXIMUM MARKS: 100

- NOTE:**
- (i) Candidate must write **Q. No.** in the **Answer Book** in accordance with **Q. No.** in the **Q. Paper.**
 - (ii) Attempt **FIVE** questions in all by selecting **TWO** questions from **SECTION-A** and **ONE** question from **SECTION-B** and **TWO** questions from **SECTION-C.** **ALL** questions carry **EQUAL** marks.
 - (iii) Extra attempt of any question or any part of the attempted question will not be considered.
 - (iv) **Use of Scientific Calculator is allowed.**

SECTION-A

Q. 1. Solve the following differential equations:

(a) $y''' - 3y'' + 2y' = \frac{e^x}{1 + e^{-x}}$ (10)

(b) $y' = \frac{2xye^{(x/y)^2}}{y^2 + y^2e^{(x/y)^2} + 2x^2e^{(x/y)^2}}$ (10)

Q. 2. (a) Find the series solution of the following differential equation:

$y'' - xy = 0$ (10)

(b) Use the method of Fourier integrals to find the solution of initial value problem with the partial differential equation.

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \quad ; \quad (-\infty < x < \infty)$$

And with initial condition $u(x, 0) = f(x)$ (10)

Q. 3. (a) Solve $x^2y'' - 3xy' + 5y = x^2 \sin(\ln x)$ (10)

(b) Find the solution of wave equation

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \text{ with boundary and initial conditions}$$

$u(0, t) = u(l, t) = 0, \quad u(x, 0) = f(x), \quad \frac{\partial u(x, t)}{\partial t} = g(x)$ (10)

SECTION-B

Q. 4. Discuss the following terms:

(5x4=20)

- (i) Tensors (ii) Kronecker delta
- (iii) Contraction (iv) Metric Tensor
- (v) Contravariant tensor of order two

Q. 5. (a) Prove that $\left\{ \begin{smallmatrix} i \\ ij \end{smallmatrix} \right\} = \frac{\partial}{\partial x^i} (\log \sqrt{g})$ (10)

(b) Prove that $\Delta = \begin{vmatrix} \delta_{m1} & \delta_{m2} & \delta_{m3} \\ \delta_{n1} & \delta_{n2} & \delta_{n3} \\ \delta_{p1} & \delta_{p2} & \delta_{p3} \end{vmatrix} = \epsilon_{mnp}$ and $\epsilon_{ijk} \epsilon_{mnp} = \begin{vmatrix} \delta_{mi} & \delta_{mj} & \delta_{mk} \\ \delta_{ni} & \delta_{nj} & \delta_{nk} \\ \delta_{pi} & \delta_{pj} & \delta_{pk} \end{vmatrix}$

Hence prove that $\epsilon_{ijk} \epsilon_{mnp} = \delta_{im} \delta_{jn} - \delta_{in} \delta_{jm}$ (10)

APPLIED MATHS, PAPER-II

SECTION-C

- Q. 6.

(a)

(i)

What is the difference between secant and false position method?
Show also graphically.

(ii)

Prove that

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

(b)

Solve the following system by Jacobi method. (Up to four decimal places).

$$\begin{aligned} 8x + y - z &= 8 \\ 2x + y + 9z &= 12 \\ x - 8y + 12z &= 35 \end{aligned}$$

(5+5=10)

Q. 7.

(a)

Evaluate by $\frac{3}{8}$ Simpson's rule

$$\int_0^3 x\sqrt{1+x^2} dx$$

; with n = 6

Also calculate the absolute error.

(b)

The amount A of a substance remaining in a reacting system after an interval of time t in a certain chemical experiment is given by following data:

A:	94.8	87.9	81.3	68.7
t:	2	5	8	14

Find t when A=80.

(10)

Q. 8.

(a)

If $f(x) = x^3$, show that $f(a,b,c) = a + b + c$

(b)

Solve by trapezoidal rule

$$\int_0^{2\pi} x \sin x dx$$

; with n = 8

(10)

<https://csshub.com.pk/>